

Biocontrol potential of *Cotesia ruficrus* Hal. (Hymenoptera : Braconidae) against different Lepidopterous pests

S.S. Patil¹, Chandani Kamble² and T.V. Sathe³

¹Dept. of Zoology, Krishna Mahavidyalaya, Rethare (Bk.) Karad, Dist. Satara.
^{2,3}Dept. of Zoology, Shivaji University, Kolhapur - 416 004, India

*Email: profdrtsathe@rediffmail.com

ABSTRACT

Cotesia ruficrus Hal. (Hymenoptera : Braconidae) is internal, larval, gregarious and polyphagous parasitoid. Wide host range is desirable attribute of an ideal parasitoid. *Heliocoverpa armigera* (Hubner), *Mythimna separata* Walker, *Agrotis ipsilon* (Hufnagel), *Spodoptera litura* Fab., *S. exigua* (Hubner) and *Exelastis atomosa* Walsingham have been tried for host preference and biocontrol potential of *C. ruficrus*. The order of preference was *P. separata* > *H. armigera* > *A. ipsilon* > *S. litura* > *S. exigua*. The most suitable host for maximum progeny production (55.00) was *M. separata*.

Keywords: *Cotesia ruficrus*, biocontrol potential, pest insects.

INTRODUCTION

Cotesia ruficrus Hal. (Hymenoptera : Braconidae) is endolarval, gregarious and polyphagous parasitoid which acts as good biocontrol agent of Lepidopterous insect pests such as *Mythimna separata* Walker, *Helicoverpa armigera* (Hubner), *Triplusia ni* Hubner, *Pseudoplusia includens* (Wlk.), *Agrotis ipsilon* Rottenberg, *Spodoptera litura* Fab, *S. exigua* (Hubner), *S. exempta* (Walker), *Sesamia cretica* Lederer, *Euro spinifera* Hubner, *Leucania loreyi*, *Plusia* spp., etc. It is distributed in Afrotropical region, Cameron, Madagascar, Senegal, Somalia, South Africa, Sudan, Uganda, Oriental and Palaearctic regions. Identifying natural enemies and their potentials on control agents is the first step in the development of biological control of insect pests. Parasitic hymenoptera discloses many of the factors that determine whether or not any two given species are to be associated as host and a parasitoid (Vinson, 1976).

How to Site This Article:

S.S. Patil, Chandani Kamble and T.V. Sathe. 2016. Biocontrol potential of *Cotesia ruficrus* Hal. (Hymenoptera: Braconidae) against different Lepidopterous pests. *Biolife*. 4(2), pp 343-346.
doi: <https://dx.doi.org/10.5281/zenodo.7318041>

Received: 5 April 2016;

Accepted; 24 May 2016; Available online : 6 June 2016

A parasitoid limits its attack to a fraction of the suitable host species that occur in nature, has stimulated investigators to study the behavior of host specificity and host selection patterns in parasitoid host interactions (Sathe, 2014). The host preference in parasitoids is vary with the species (Weshloh, 1976). Wide host range is desirable attribute of an ideal parasitoid. The parasitoids can breed in the laboratory on unnatural hosts, is often of great importance in the mass propagation of parasitoids, their field release against target pest species and their colonization. Therefore, the objective of the present work was to search for new hosts and biocontrol potential on the different insect pests. Review of literature indicates that Hafez (1947), Broodryk (1969), Fisher (1971), Odebiyi and Oatman (1972), Vinson and Guillot (1972), Jackson *et al.* (1979), Hopper and King (1984), Sathe (1987, 1988, 1990), Sathe & Santhakumar (1992), Sathe & Margaj (2001), Sathe *et al.*, (2003), Sathe & Oulkar (2010), Sathe (2014), Sathe & Chougale (2014), Sutar and Sathe (2016), Jadhav & Sathe (2016) etc. attempted the studies related to parasitoid host preferences.

MATERIAL AND METHODS

Culture of *C. ruficrus* and its insect hosts *H. armigera*, *A. ipsilon*, *M. separata*, *S. litura*, *S. exigua* and *Exelastis atomosa* Walsingham were started in the laboratory (27±1°C, 75-80%, 12 hr photoperiod) by

providing respective hosts and host food plants parts to parasitoid and hosts respectively. 6-7 day old larvae of above pests with 20 density were exposed to mated female parasitoid in a glass cage (25 x 25 x 25 cm) for 2 hr for oviposition. The parasitized hosts were reared on their respective host plants in plastic containers individually for screening parasitoids. Records were made on per cent mortality of host species, per cent host emergence and per cent parasitism by *C. ruficrus* to different insect pests used in the experiments. All experiments were replicated for five times for confirmation of results.

RESULTS

Results are recorded in table - 1 and figs 1-4. The results indicated that *C. ruficrus* preferred the host *M. separata* at its maximum biocontrol potential (parasitism) of 55.00%. *H. armigera*, *A. ipsilon*, *S. litura*, *S. exigua* and *E. atomosa* showed 50.00%, 45.00%, 40.00%, 25.00% and 0.00% biocontrol potential respectively.

The order of host preference given by *C. ruficrus* was *M. separata* > *H. armigera* > *A. ipsilon* > *S. litura* > *S. exigua*. The larvae of *E. atomosa* were remained unparasitized by *C. ruficrus*. The host mortality rate was 00.00%, moth emergence was 35.00% and percent parasitism by *C. ruficrus* to *M. separata* was 55.00%. *H. armigera* was also found to be potential host for production of progeny of parasitoid by 50.00% parasitism (Table-1).

DISCUSSION

Biology *C. ruficrus* was studied by McCutcheon *et al.* (1983) by providing hosts *Pseudoplusia includens*, *T. ni* and *Spodoptera frugiperda*. The development from egg to adult was possible within the range of 16 to 18 days on above said hosts and mated females survived for 18 days and unmated ones for 20 days. Hafez (1947) also studied the biology of *C. ruficrus* by providing *A. ipsilon* larvae wherein the parasitoid completed its life cycle from egg to adult within 14 to 24 days while, in the present study, the life cycle on *H. armigera* was

Table-1. Host preference and biocontrol potential of *C. ruficrus* against some lepidopterous pests

Sr. No.	Host species	Host density	% Host mortality	% moth emergence	% parasitism
1	<i>H. armigera</i>	20	10.00	40.00	50.00
2	<i>M. separata</i>	20	00.00	35.00	55.00
3	<i>A. ipsilon</i>	20	10.00	45.00	45.00
4	<i>S. litura</i>	20	15.00	45.00	40.00
5	<i>S. exigua</i>	20	15.00	60.00	25.00
6	<i>E. atomosa</i>	20	10.00	90.00	00.00



Fig. 1: *C. ruficrus* (Adults)



Fig. 2: *C. ruficrus* (Cocoons)



Fig. 3: *M. separata* (Larvae)



Fig. 4: *A. ipsilon* (Larva)

completed within 16-18 days by the parasitoid under laboratory conditions ($27\pm 1^{\circ}\text{C}$, 75-80%, 12 hr photoperiod).

For successful parasitism, discrimination of host suitability for progeny often follows host recognition. Through a long history of random observations, receptors on the antennae, tarsi and ovipositor have been implicated in this phase of a parasitoid's behaviour (Fisher, 1971). Some braconids can distinguish host suitability, other species reportedly cannot (Vinson & Guillot, 1972; Sathe, 1988).

According to Broodryk (1969) *Chelonus* (*Microchelonus*) *curvimaculatus* Cameron parasitized the lepidopteran hosts such as *Phthorimaca operculella* (Zeller), *Cryptophebia leucotreta* (Meyr.) *Ephestia cautella* (Wlk.), *E. kuchniella* (Zeller), *S. exempta*, *S. exigua*, *S. litura* and *H. armigera*. However, *Bombyx mori* L. was found unsuitable for parasitism. In the present study, the hosts belong to the family Noctuidae were parasitized but, *E. atomosa* which was from the family Pterophoridae was not parasitized by the parasitoid.

Agathis gibbosa (Say) (Braconidae) showed increased number of progeny production with respect to host density 25, 50, 75, 85, 100 and 125 but the number of parasitoid was decreased after host density 150. The host species exposed by Odebiyi & Oatman (1972) was *P. operculella*.

Hopper and King (1984) studied the preference of *Microplitis croceipes* (Cresson) (Hymenoptera: Braconidae) towards the instars of *Heliothis zea* (Bodalic) and *H. virescens* (F.) on cotton crop. The wasp preferred third instar larvae of host species most. The order of preference of instars by *M. croceipes* was third instars > fourth instar > second instars. However, the sex ratio of parasitoid was not affected significantly by the instar parasitized.

In an Ichneumonid *Campoletis chlorideae* (Uchida) (Hymenoptera : Ichneumonidae), an internal, larval parasitoid of *H. armigera* host preference was studied by Sathe & Santhakumar (1992) by providing *H. armigera*, *S. litura*, *S. exigua* and *E. atomosa*. Wherein *E. atomosa* larvae were rejected by the parasitoid. The order of preference shown by the parasitoid was *H. armigera* > *S. litura* > *S. exigua* with 30.4%, 21.6% and 17.6% parasitism respectively.

According to Sathe & Margaj (2001) *A. arterus* gave preference to only *Erias insulana* while, *A. pusaensis* gave preference to only *Sylepta derogata* when exposed the hosts, *H. armigera*, *P. gossypiella*, *S. litura* and *S. derogata*.

According to Jackson *et al.* (1979) suitable hosts for *Chelonus* (*Microchelonus*) *blackburni* Cameron were *Pectinophora gossypiella* (Saunders), *Heliothis zea* (Boddie), *H. virescens* (F.), *T. ni* and *S. exigua*. However, Salt marsh caterpillars *Estigmene acrea* (Drum) were not accepted for parasitism by *C. blackburni*. A dipterous parasitoid *Exorista bombycis* L. gave more preference to *Bombyx mori* L. Fc_2 pure line breed than Fc_1 (Jadhav & Sathe, 2016).

Investigation of different hosts for biocontrol agents is a challenging aspect and is the need of the day for strengthening biological pest control programmes as an ecofriendly tool of pest management. The present work will add great relevance for strengthening biological pest control strategies.

ACKNOWLEDGEMENTS

Authors are thankful to Shivaji University, Kolhapur and Krishna Mahavidyalaya, Rethare (Bk.) Karad, Dist. Satara for providing facilities to this work.

Conflict of Interests

Authors declare that there is no conflict of interests regarding the publication of this paper.

REFERENCES

- [1]. Fisher, R.C. Aspects of Physiology of endoparasitic hymenoptera. *Biol. Res.* **46**, 243-278 (1971).
- [2]. Hopper, K.R. and E.G. King. Preference of *Microplitis croceipes* (Hymenoptera : Braconidae) for instars and species for *Heliothis* (Lepidoptera : Noctuidae). *Environ. Entomol.*, **15**, 1145-1150 (1984).
- [3]. Jackson, D.J., Neemann, E.G. and R. Patana. Parasitization of six lepidopteran cotton pests by *Chelonus blackburni* (Hymenoptera : Braconidae). *Entomophaga*, **24**, 99-105 (1979).
- [4]. Jadhav A.D. and T.V. Sathe. Host preference by Uzi fly *Exorista bombycis* L. in pure line bivoltine breeds Fc_1 and Fc_2 (*Bombyx mori* L.) and economical loss in seed cocoon production. *Biolife*, **4**(1), 88-93 (2016).
- [5]. Odebiyi J.A. and Oatman E.R. Biology of *Agathis gibbosa* (Hymenoptera : Braconidae), a primary parasite of the potato tuberworm. *Ann. ent. Soc. Am.* **65**, 1104-1114 (1972).
- [6]. Sathe T.V. New records of parasitoids of Ber hairy caterpillar *Thiocidas postica* Wlk. in Kolhapur, India. *Sci. & Cult.* **53**, 185 (1987).
- [7]. Sathe T.V. Impact of host food plant on parasitization behaviour in a larval parasitoid of *Heliothis armigera* (Hubn.) *Proc. Indian Acad. Sci. (Anim. Sci.)*, **99**, 233-242 (1990).
- [8]. Sathe T.V. Recent trends in Biological pest control. Astral International Pvt. Ltd., New Delhi pp 204 (2014).
- [9]. Sathe T.V. Biological control through Ichneumonids. Astral International Pvt. Ltd., New Delhi. pp. 1-117 (2015).
- [10]. Sathe T.V. & T.M. Chougule. Hymenopterous biopesticides and their preliminary biocontrol potential from Western Maharashtra including Ghats. *Biolife*, **2**(4), 1254-1261 (2014).

- [11]. Sathe T.V. and S.G. Margaj. Cotton pests and biocontrol agents. Daya Publishing House, New Delhi. pp. 1-166.
- [12]. Sathe T.V. and Jyoti M. Oulkar. Insect pest management - Ecological concepts. Daya Publishing House, New Delhi. pp. 1-235 (2010).
- [13]. Sathe T.V. and M.V. Santhakumar. Host specificity in *Campoletis chloridae* (Uchida) (Ichneumonidae : Hymenoptera). *J. Adv. Zool.*, **13(1 & 2)**, 53-56 (1992).
- [14]. Sathe, T.V., S.A. Inamdar and R. Dawale. Indian pest parasitoids. Daya Publishing House, New Delhi. pp. 1-145 (2003).
- [15]. Sutar Mahesh and T.V. Sathe. Diversity and biocontrol potential of the genus *Diadegma* Cameron (Hymenoptera : Ichneumonidae) from *Biolife*, **4(1)**, 202-208 (2016).
- [16]. Thurston, R. and Postley, L. Effect of instars of *Manduca sexta* on the rate of development of its parasite *Apanteles congregatus*. *Tob. Sci.*, **22**, 32-34 (1978).
- [17]. Vinson S.B. Host selection by insect parasitoids. *A. Rev. Ent.*, **21**, 109-134 (1976).
- [18]. Vinson S.B. and Guillot F.S. Host marking source of a substance that results in host discrimination in insect parasitoids. *Entomophaga*, **17**, 241-245 (1972).
- [19]. Weshloh. Behavioural responses of the parasite *Apanteles melanoscelus* to gypsy moth silk. *Environ. Entomol.*, **5**, 1128-1132 (1976).

DOI: