Biolife
AN INTERNATIONAL QUARTERLY JOURNAL OF BIOLOGY & LIFE SCIENCES
2(4):1310-1315
ISSN (online): 2320-4257
www.biolifejournals.com
BIOLIFE RESEARCHARTICLE

SOD and CAT level in the abdominal muscles of immunostimulated mice during hepatitis B infection

Madhuri, D[1] and Viveka Vardhani,V[2]*

[1] *Department of Biochemistry, AcharyaNagarjuna University, Nagarjunanagar-522 510.*
[2]*Department of Zoology, AcharyaNagarjuna University, Nagarjunanagar-522 510.*

E-mail: vadlamudi_vv@yahoo.co.in

ABSTRACT

The effect of immunostimulant, IDS and Gen Vac B vaccine was studied on the enzymatic activities of SOD and CAT in the abdominal muscles of male Swiss albino mice mice (6 - 8 weeks old; 23 - 26g wt) at different days of experiment. Eighty mice were separated into 8 groups as group I (Immunex DS was orally administered @ 150mg/mouse on 0 day), group A (IDS @ 150 mg/mouse + 0.07 ml/mouse Gen Vac B vaccine was injected on day 4 of experiment, group B (IDS @ 150mg/mouse + vaccine @ 0.1 ml/mouse), group C (IDS @ 150mg/mouse + vaccine @ 0.2 ml/mouse), group D (IDS @ 150mg/mouse + vaccine @ 0.4 ml/mouse), group E (IDS @ 150mg/mouse + vaccine @ 0.8 ml/mouse) and group F (IDS @ 150mg/mouse + vaccine @ 1.0 ml/mouse) and group U (untreated with IDS + uninfected) was kept as controls for comparison. Two mice from each experimental (A, B, C, D, E and F) and control groups U and I (after day 7 of vaccine treatment in case of experimentals) were necropsied, from day 1-5 of experiment. Abdominal muscle tissue was separated and analyzed for SOD and CAT activities using standard methods. The activity of SOD showed a considerable increase in all the experimental groups of mice and a decreased CAT activity when compared with controls (group U) and IDS treated mice (group I) during the entire experimental period. The level of SOD and CAT almost remained constant from day 1 to 5 of experiment in IDS treated mice. Increased activities of these of enzymes are responsible for regulation of active oxygen species which play a significant role in pathogenesis of muscular dystrophies. It is evident that IDS and/or vaccine might have caused stress resulting in the marked alteration in the level of SOD and CAT in the abdominal muscles of mice.

Keywords: SOD, CAT, Abdominal muscles, Mice, Immunostimulant, Hepatitis B.

INTRODUCTION

More than 300 million people worldwide are surviving as hepatitis B Virus (HBV) carriers (Lavanchy, 2004; Milich and Liang 2003). Hepatitis B is the common type of cancer which is highly reported in Asians, particularly Chinese and Indians (Wong and Goh, 2006). One way to suppress inflammatory and/or stimulation of phagocytosis is the application of immunostimulants which increase resistance to bacterial and viral infections by stimulating nonspecific immune mechanisms in mice (Petrunov et al., 2007). Certain antioxidants protect the body against the damages caused by ROS (destructive oxygen derivatives) and maintain redox homeostasis; these include both enzymatic (SOD), glutathione peroxidase (GPx), catalase (CAT), etc and non-enzymatic

(glutathione and vitamins A, E and C) antioxidants. Oxidative stress sets in when the redox balance is disrupted by extreme generation of ROS or when the antioxidant capacity is insufficient (Thomas, 2000; Golden et al., 2002). The role of oxidative stress in the mechanism of isoniazid and rifampicin-induced hepatitis has been reported by Attri et al., (2000). SOD and CAT play an important role in the biological systems to act against oxidative stress (Akyol et al., 2002). Oxygen is required to maintain life and metabolic processes including drug metabolism but ROS are generated during oxygen use (Gupta et al., 2007). It was observed that the bark extract (*Helicteresisora*)could increase the SOD and CAT activities in the cardiac tissues of diabetic rats (Kumar et al., 2008). SOD activity was decreased in both liver and kidney whereas CAT activity was increased only in liver in cadmium exposed rats and reversed on selenium administration (Ognjanovic et al., 2008). Hao et al., (2009) reported that purslane*(Portulacaoleracea)* can be used for anti-aging, thereby increasing the level of SOD and decreasing the level of MDA in the brain of mice treated with D-galactosamine. GSH homeostasis in tissues is maintained by the antioxidant enzymes such as SOD and GST (Abdel-Moneim*et al*., 2010). The hepatic antioxidant enzymatic (SOD and CAT) activites were decreased in the liver of rats administered with CCl_4, and restored by *Decalepishamiltonii* (DHA treatment) (Srivastava and Shivanandappa, 2010). Significant acceleration in the level of lipid peroxide (MDA), with significant depletion in GSH, SOD and CAT activities in the serum of rats exposed to two different doses of gamma radiation that produced oxidative stress (Saad and Ammar, 2011). There was a significant increase of liver, kidney and testes catalase and SOD activities in purslaneadministered albino rats (Dkhil et al., 2011). D-galactosamine/ lipopolysaccharide (D-GalN/ LPS) intoxicated rats showed a decreased activity of hepatic antioxidants like SOD and CAT (Fyiad et al., 2012). Decreased activities of hepatic antioxidant enzymes, such as SOD and CAT were observed in isoniazide and rifampicin treated rats (Verma et al., 2013). Investigations on the hepatoprotective effects of Triphala in D-Galactosamine (D-GalN) induced hepatic toxicity in mice explain that D-GalN induced hepatic damage resulted in a significant increase in the activity of ALT, AST, ALP, bilirubin, lipid peroxidation (LPO) and Tumour necrosis factor (TNF-α) level and a decrease in the levels of anti-oxidant enzymes such as SOD, CAT, glutathione peroxidase (GPx), glutathione reductase (GR), glutathione-s-transferase (GST) and total reduced glutathione (Sabina et al., 2013). The activities of SOD and CAT were significantly lowered in tissues (liver, proximal colon, distal colon and lysate) of 1, 2-dimethylhydrazine (DMH) treated rats and linalool supplementation significantly increased the activities of SOD and CAT (Srithar*et al*., 2013). Pretreatment with *Tribulusterrestris*fruit aqueous extract (TTFAEt) maintained the normal endogeneous antioxidant activity by increasing the levels of antioxidant enzymes such as SOD and CAT in heart tissue of rats (Sailaja*et al*., 2013). Oils from *Zinger officinale*and *Curcuma longa* and at a dose of 200mg/kg showed hepatoprotection by restoring the activitiy of SOD in ethanol-treated rats (Nwozo et al., 2014).Decreased activities of CAT and SOD was observed in all the tissues of brain, liver and kidney of mice exposed to sodium fluoride (NaF) compared to control (Sandeep et al., 2014). The present investigations are carried out to understand the influence of IDS in SOD and CAT level in mice treated with Hepatitis B infection

MATERIAL AND METHODS

According to the guidelines of CPCSEA, eight groups (10 in each group) of male Swiss albino mice (*Musmusculusalbinus*) (6 - 8 weeks old; 23 - 26g wt) were fed with standard balanced diet and water *ad libitum*and taken care. Immunex DS (IDS) was orally administered @150mg/mouse to group I (10 mice). Another 6 groups of mice received IDS orally @150mg/mouse on 0 day and Gen Vac B Vaccine on day 4 of experiment; @ 0.07 ml/mouse in group A, IDS @ 150mg/mouse + 0.1ml/mouse in group B, IDS @ 150mg/mouse +

Table-1:Superoxide dismutase (units/mg of protein/min) a abdominal muscles of experimental (Group A - treated with with Hbs Ag @ 0.07 ml/mouse), (Group B - treated with Imm Hbs Ag @ 0.1 ml/mouse), (Group C - treated with Immunex D @ 0.2 ml/mouse), (Group D - treated with Immunex DS (@ 15 ml/mouse), (Group E - treated with Immunex DS @ 150 m ml/mouse), (Group F - treated with Immunex DS @ 150 mg/mo

Table -2 : t values obtained in SOD levels of different experimental groups (A, B, C, D, E and F) of mice.

	Experimental groups						Control groups	
SOD :	**A**	**B**	**C**	**D**	**E**	**F**	**U**	**I**
Mean	2.75	3.28	4.30	5.91	7.95	12.13	1.89	2.87
t values	A–U t = 2.21@		B–U t = 4.49*		C–U t = 3.96*			
	D–U t = 3.19*		E–U t = 3.41*		F–U t = 3.05*			
	A–I t = 0.29@		B–I t = 1.32@		C–I t = 2.33*		I–U t = 17.40*	
	D–I t = 2.41*		E–I t = 2.85*		F–I t = 2.76*			
	A–B t = 1.07@		A–C t = 2.14@		A–D t = 2.38*		A–E t = 2.86*	A–F t = 2.78*
	B–C t = 1.47@		B–D t = 2.02@		B–E t = 2.58*		B–F t = 2.62*	
	C–D t = 1.14@		C–E t = 1.92@		C–F t = 2.30@			
	D–E t = 0.93@		D–F t = 1.75@					
	E–F t = 1.10@							

P value at 5% level of significance is 2.306. - Statistically significant values. @ - Statistically non – significant values.*

0.8ml/mouse in group E and1ml/mouse in group F, Another group (U) of ten mice was kept as controls for comparison (untreated with IDS + uninfected). Two mice from each of the experimental (after day 7 of vaccine treatment) and control groups were sacrificed on day 1, 2, 3, 4 and 5 of experiment, abdominal muscle tissue was separated and analysed for the activities of SOD and CAT following the methods of Misra and Fridovich (1972) and Forman and Fischer (1981). Results were analysed for statistical significance using student's t test.

RESULTS AND DISCUSSION

The activity of SOD (table 1) showed a considerable increase in all the experimental groups of mice when compared with controls and IDS treated mice (except the level of SOD on day 1 and 2 in groups A and B). Whereas, a decreased CAT activity was found during the entire experimental period (except from day 3 to 5 in groups A, C and on day 4 and 5 in group D and on day 5 in group F) when compared with controls (group U) and with immunostimulated

Table 3 : t values obtained in CAT levels of different experimental groups (A,B,C,D,E and F) of mice

	Experimental groups						Control groups	
CAT :	**A**	**B**	**C**	**D**	**E**	**F**	**U**	**I**
Mean	15.36	9.58	16.73	13.51	6.29	9.15	12.87	13.86
t values	A–U t = 1.36@	B–U t = 5.22*	C–U t = 1.17@					
	D–U t = 0.29@	E–U t = 7.17*	F–U t = 2.63*					
	A–I t = 0.82@	B–I t = 6.78*	C–I t = 0.87@				I–U t =17.99*	
	D–I t = 0.16@	E–I t = 8.25*	F–I t = 3.33*					
	A–B t = 3.02*	A–C t = 0.36@	A–D t = 0.65@				A–E t = 4.43*	A–F t = 2.68*
	B–C t = 2.18@	B–D t = 1.74@	B–E t = 2.89*				B–F t = 0.24@	
	C–D t = 0.90@	C–E t = 3.06*	C–F t = 2.12@					
	D–E t = 3.07*	D–F t = 1.68@						
	E–F t = 1.69@							

P value at 5% level of significance is 2.306. - Statistically significant values. @ - Statistically non – significant values.*

mice (group I) (except on day 4 and 5 in groups A, D and from days 3 to 5 in group C and on day 5 in F).

There was a significant difference in the level of SOD in all the experimental groups (except in group A) when compared with controls (group U) and with immunostimulated mice (group I) (except in groups A and B). There was no significant difference in the level of SOD in all the experimental groups (except in groups A with D, E and F and B with E and F) when compared among themselves (Table 2). There was a significant increase in the level of CAT in groups B, E and F and a significant decrease in groups A , C and D when compared with controls (group U) and when compared with IDS treated mice there was a significant difference in all the groups (except A, C and D). There was no significant diference in the level of CAT in all the experimental groups (except in groups A with B, E and F, B with E and C, D with E) when compared among themselves (Table 3).

The present investigations indicate that vaccination might have brought intolerance of some inherited factors like glycogenolytic defect in the muscle, though the role of these factors in the activation of enzymes in muscles is not known, Also, it is clear that muscle fibers changed their properties during new activities as stated by Pette and Vrbova (1985).

ACKNOWLEDGEMENTS

The author (Madhuri, D) is thankfull to UGC, New Delhi for financial assistance in the form of RGNF and to Prof. PVV Satyanarayana, the then Head of the Department of Biochemistry for providing laboratory facilities.

REFERENCES

1. Abdel-Moneim, AE.,Dkhil, MA. AI-Quraishy, S. 2010. The redox status in rats treated with flaxseed oil and lead-induced hepatotoxicity. Biol. Trace Elem. Res. (Epub ahead of print).
2. Akyol, O., Herken, H., Uz, E., Fadillioglu, E., Unal, S., Sogut, S., Ozyurt, H. and Savas, HA. 2002. The indices of endogenous oxidative and antioxidative processes in plasma from schizophrenic patients. The possible role of oxidant/antioxidant imbalance. Prog. Neuropsychopharmacol Biol. Psychiarty. 26: 995-1005.
3. Attri, S., Rana, S. V., Vaiphei, K., Sodhi, CP., Katyal, R., Goel, RC et al. 2000. Isoniazid and rifampicin induced oxidative hepatic injury protection by N-acetycysteine. Hum Exp. Toxicol.19: 517-522.
4. Dkhil, M. A., AbdelMoniem, A E., AI-Quraishy, S. and Saleh, R. A. 2011. Antioxidant effects of purslane (Portulacaoleracea) and its mechanism of action. J. Medicinal. Plants. Research. 5(9): 1589-1563.
5. Forman, H.J. and Fischer, A. B. 1981 Antioxidant defense in: Danial, E and Gilbert, L: eds. Oxygen and living process. An inter disciplinary approaches. Academic Press, New York 237-248.
6. Fyiad, A. A., Abd El-Kader, M. A. and Abd El-Haleem, A. H. 2012. Modulatory effects of pomegranate juice on nucleic acids alterations and oxidative stress in experimentally hepatitis rats. Life Science J. 9(3): 676-682.
7. Golden, T. R., Hinerfeld, D. A. and Melov, S. 2002. Oxidative stress and aging: Beyond correlation. Aging Cell. 1: 117-123.
8. Gupta, S., Agarwal, A., Banerjee, J. and Alvarez, J. G. 2007. The role of oxidative stress in spontaneous abortion and recurrent pregnancy loss: A systemic review. Obstet. Gynecol. Surv. 62: 335-347.
9. Hao, H., Nancai, Y., lei, F., Wen, S., Guofu, H., Yanxia, W., Hanju, H. and Qian, L.2009. Retracted: Antiaging effect of purslane herb aqueous extracts and its mechanism of action. Phytother. Res. 23: i-vii.
10. Kumar, G., Banu, G. S. and Murugesan, A. G. 2008. Effect of *Helicteresisora* bark extracts on heart antioxidant status and lipid peroxidation in streptozotocin diabetic rats. J. Appl. Biomed. 6: 89-95.
11. Lavanchy, D. 2004.Hepatitis B virus epidemiology, disease burden, treatment and current and emerging prevention and control measures. J. Viral Hepatol. 11(2): 97-107.

12. Milich, D. and Liang, J. T. 2003. Exploring the Biological basis of hepatitis B e antigen in hepatitis B virus infection. Hepatol. 38(5): 1075-1086.
13. Misra, H. P. and Fridovich, I. 1972. The role of superoxide anion in the antioxidation of epinephrine and a sample assay for superoxide dismutase. J. Biol. Chem. 247: 3170-3175.
14. Nwozo, S. O., Osunmadewa, D. A. and Oyinloye, B. E. 2014. Anti-fatty liver effects of oils from *Zinger officinale*and *Curcuma longa* on ethanol-induced fatty liver in rats. J. Integr. Med. 12(1): 59-65.
15. Ognjanovic, B. I., Markovic, S. D., Pavlovic, S. Z., Zikic, R. V. and Stajn, A. S. 2008. Effect of chronic cadmium exposure on antioxidant defense system in some tissues of rats: protective effect of selenium. Physiological Research 57(3): 403-411.
16. Petrunov, B., Nenkov, P. and Shekerdjiisky, R. 2007. The role of immunostimulants in immunotherapy and immunoprophylaxis. Biotechnol. Biotechnol. EQ. 21(4): 454-462.
17. Pette, D and Vrbova, G. 1985. Neural control of phenotypic expression in mammalian muscle fibers. Muscle Nerve. 8: 676-689.
18. Saad, TM. M. and Ammar, AA. A. 2011. Protective role of peanut oil in rats exposed to two different doses of gamma radiation that produced oxidative stress and bone injury. Egyptian. J. Hospital Medicine. 44: 380-391.
19. Sabina, E. P., Rasool, M., Vedi, M. and Geethanjali, A. 2013.Protective properties of traditional herbal formulation against D-Galactosamine induced hepatotoxicity in mice. Int. J. Drug Dev. & Res. 5(2): 164-173.
20. Sailaja, K.V., Shivaranjani, V. L., Poornima, H., Md, S. B., Rahamathulla, K., Devi, L* 2013. Protective effect of *TribulusTerrestris* L. fruit aqueous extract on lipid profile and oxidative stress in isoproterenol induced myocardial necrosis in male albino wistar rats. EXCLI J. 12: 373-383.
21. Sandeep, V., Kavitha, N., Praveena, M., Sekhar, P. R. and Rao, K. J. 2014. Alterations of detoxification enzyme levels in different tissues of sodium fluoride (NaF) treated albino mice. Int. J. Adv. Research. 2(1): 492-497.
22. Srithar, G., Sudha, M. and Nalini, N. 2013. Linalool exerts dose dependent chemopreventive effect against 1,2-dimethyhydrazine induced rat colon carcinogenesis. Int. J. Pharmceu. and Biol. Archives. 4(4): 758-770.
23. Srivastava, A. and Shivanandappa, T. 2010. Hepatoprotective effect of the root extract of *Decalepishamiltonii* against carbon tetrachloride-induced oxidative stress in rats. Food Chemistry. 118: 411-417.
24. Thomas, M. J. 2000. The role of free radicals and antioxidants. Nutrition. 16: 716-718.
25. Verma, V. K. and Hussain, Z. 2013. Effect of 50% ethanolic extract of roots of *Cissampelospareira* on some bio-chemical parameters to investigate the hepato-protective study in animal models (rats). J. Drug Discovery and Therapeutics. 1(7): 106-116.
26. Wong, C. H. and Goh, K. L. 2006. Chronic hepatitis B infection and liver cancer. Biomed. Imaging. Interv J. 2(3): e7.

DOI:
https://dx.doi.org/ 10.5281/zenodo.7238924
Received: 9 October 2014;
Accepted; 25 November 2014;
Available online : 14 December 2014